

CORRECTION

Open Access



Correction to: The elevated levels of urinary angiotensinogen are correlated with the severity of idiopathic membranous nephropathy

Ziyong Tang¹, Yue Wang¹, Liyuan Tao², Yanhong Guo¹, Yimu Zheng¹ and Danxia Zheng^{1*}

Correction to: *BMC Nephrology* (2018) 19:357
<https://doi.org/10.1186/s12882-018-1165-1>

Following publication of the original article [1], it was reported that, due to a typesetting mistake, the three tables and two figures for this article were included as an Additional file instead of in the body of the article. The original publication of this article has been updated to correct this.

The publisher apologises to the authors and readers for the inconvenience.

Author details

¹Department of Nephrology, Peking University Third Hospital, No. 49, North Garden Road, Haidian District, Beijing 100191, China. ²Department of Clinical Epidemiology, Peking University Third Hospital, Beijing, China.

Received: 13 December 2018 Accepted: 13 December 2018

Published online: 13 February 2019

Reference

1. Tang, et al. The elevated levels of urinary angiotensinogen are correlated with the severity of idiopathic membranous nephropathy. *BMC Nephrol.* 2018;19(357). <https://doi.org/10.1186/s12882-018-1165-1>.

* Correspondence: dxzheng@sina.com

¹Department of Nephrology, Peking University Third Hospital, No. 49, North Garden Road, Haidian District, Beijing 100191, China

Full list of author information is available at the end of the article

